

# Big mapping class groups and rigidity of the simple circle

Lvzhou Chen  
joint work with Danny Calegari


Hyperbolic Lunch, UToronto  
June 3, 2020

# Big MCG and actions on $S^1$

$\Omega = \mathbb{R}^2 - K$ ,  $K$  Cantor set,  $\Gamma = \text{Mod}(\Omega)$ .

**Fact:**  $\Gamma$  acts **faithfully** on  $S^1$ .

$\implies \Gamma \hookrightarrow \text{Homeo}^+(S^1)$ .



**Question 1:** Does  $\Gamma$  act on  $S^1$  in different ways?

Any nontrivial action with a global fixed point?

**Question 2:** Is  $\Gamma$  generated by torsion?

$\text{Homeo}^+([0, 1])$  is torsion-free.

**Question 3:** Any further obstruction for  $G \hookrightarrow \Gamma$ ?

# Main results

**Question 1:** Does  $\Gamma$  act on  $S^1$  in different ways? **No.**

**Question 2:** Is  $\Gamma$  generated by torsion? **Yes.**

**Question 3:** Any further obstruction for  $G \leq \Gamma$ ? **Yes.**

**Theorem 3:** Every **countable** subgroup of  $\text{Homeo}^+(S^1)$  embeds into  $\Gamma$ . Not true if uncountable, e.g.  $\text{PSL}_2\mathbb{R}$ .

**Theorem 2:**  $\Gamma$  is normally generated by a single 2-torsion.

**Theorem 1:**  $\Gamma$  acts faithfully and **minimally** on the **simple circle**  $S_S^1$ . Any nontrivial action of  $\Gamma$  on  $S^1$  is **semi-conjugate** to this one.

$$S^1 \xrightarrow{\rho(g)} S^1$$

$$h \downarrow$$

$$h \downarrow$$

$$S_S^1 \xrightarrow{g} S_S^1$$

Similar results by Mann–Wolff

# Rays


$S$  finite type  
curve graph  $\mathcal{C}(S)$   
 $\mathcal{C}(S)$  is hyperbolic  
(Masur–Minsky)

$S = \Omega$   
ray graph  $\mathcal{R}$   
 $\mathcal{R}$  is hyperbolic  
(Bavard,  
Aramayona–Fossas–Parlier)

# Rays

Fix a hyperbolic structure on  $\Omega$ .

conical cover  $\Omega_C$  and conical circle  $S_C^1$ .



# Rays

Fix a hyperbolic structure on  $\Omega$ .

conical cover  $\Omega_C$  and conical circle  $S_C^1$ .


**Simple** rays =  $R \sqcup L \sqcup X$ ,  $\Gamma$ -invariant and closed.

$R$  = short rays,  $L$  = lassos,  $X$  = long rays.

one orbit



one orbit

uncountably many orbits, related to  $\partial\mathcal{R}$



# Unique minimal set

**Lemma:** For any  $x \in S_C^1$ , the orbit closure  $\overline{\Gamma x} \supset R$ .




# Unique minimal set

**Lemma:** The unique minimal set  $\bar{R} = R \sqcup X$  is a Cantor set on  $S_C^1$ , complementary intervals  $\leftrightarrow L$ .

**Key points:**

- each  $\ell \in L$  is isolated in the simple set.





# Unique minimal set

**Lemma:** The unique minimal set  $\bar{R} = R \sqcup X$  is a Cantor set on  $S_C^1$ , complementary intervals  $\leftrightarrow L$ .

**Key points:**

- each  $\ell \in L$  is isolated in the simple set.  $\implies R \cup X$  closed.
- the simple set is nowhere dense.  $\implies R \cup X$  nowhere dense.
- can approximate each  $x \in X$  by short rays.  
 $\implies \bar{R} = R \cup X$ , perfect.

# The simple circle

Collapse complementary intervals to points  $S_C^1 \rightsquigarrow S_S^1$   
simple circle

- $\Gamma$  acts faithfully and minimally on  $S_S^1$
  - has an uncountable orbit  $R$ .
  - has a countable orbit  $L$ .
- $\implies$  every subgroup of  $\Gamma$  has a faithful action on  $S^1$  with a countable orbit.

# $\mathrm{PSL}_2\mathbb{R}$ does not embed

**Theorem 3:** Any countable subgroup of  $\mathrm{Homeo}^+(S^1)$  embeds into  $\Gamma$ . Not true if uncountable, e.g.  $\mathrm{PSL}_2\mathbb{R}$ .

**Prop.:** Every faithful action of  $\mathrm{PSL}_2\mathbb{R}$  on  $S^1$  is standard.

**Tool:** The **bounded Euler class**  $\mathrm{eu}_b^\rho \in H_b^2(G)$ .

$$\rho : G \rightarrow \mathrm{Homeo}^+(S^1), \rightsquigarrow \mathrm{eu}_b^\rho = \rho^* \mathrm{eu}_b.$$

- (Ghys)  $\mathrm{eu}_b^\rho$  determines the action up to semi-conjugacy.
- $\mathrm{eu}_b^\rho = 0$  iff the action has a **global fixed point**.
- $H_b^2(G) \hookrightarrow H^2(G)$  if  $G$  is uniformly perfect.  
 $\mathrm{eu}_b^\rho \mapsto \mathrm{eu}^\rho, \mathrm{eu}^\rho = 0$  iff action lifts to  $\mathbb{R}$ .

# $\mathrm{PSL}_2\mathbb{R}$ does not embed

**Theorem 3:** Any countable subgroup of  $\mathrm{Homeo}^+(S^1)$  embeds into  $\Gamma$ . Not true if uncountable, e.g.  $\mathrm{PSL}_2\mathbb{R}$ .

**Prop.:** Every faithful action of  $\mathrm{PSL}_2\mathbb{R}$  on  $S^1$  is standard.

**Proof sketch:**

- any  $g \in \mathrm{PSL}_2\mathbb{R}$  is a commutator, so uniformly perfect.
- $H^2(\mathrm{PSL}_2\mathbb{R}; \mathbb{Z}) \cong \mathbb{Z}$ , generated by  $eu^{std}$ .
- $eu^\rho = \lambda \cdot eu^{std}$ ,  $\lambda = \pm 1$  (torsion+rigid subgroup).

Then show the action  $\rho$  is transitive, so no countable orbit.

# Countable subgroups

**Theorem 3:** Any countable subgroup of  $\text{Homeo}^+(S^1)$  embeds into  $\Gamma$ . Not true if uncountable, e.g.  $\text{PSL}_2\mathbb{R}$ .

**Proof:** Denjoy's blow-up construction + suspension.

# Rigidity

**Theorem 1:**  $\Gamma$  acts faithfully and minimally on the simple circle  $S^1_S$ . Any nontrivial action of  $\Gamma$  on  $S^1$  is semi-conjugate to this one (up to a change of orientation).

**Proof sketch:** Fix an action  $\rho$  without fixed points.

$r$  short ray,  $\Gamma_r := \text{Stab}(r)$ .

**Step 1:** Each  $\Gamma_r$  acts with a global fixed point.

Fix  $r_0$ , pick  $P(r_0) \in \text{Fix}(\Gamma_{r_0})$ . Let  $P(r) = \rho(g).P(r_0)$  if  $r = g.r_0$ .

$$\begin{array}{ccc} R & \longrightarrow & S^1_S & \text{fixed by } \Gamma_r = g\Gamma_{r_0}g^{-1} \\ P \downarrow & \Gamma\text{-equivariant} & & \\ & S^1 & & \end{array}$$


**Step 2:**  $P$  preserves the circular order and is injective.

# Step 1 details

$r$  short ray,  $\Gamma_r := \text{Stab}(r)$ ,  $\Gamma_{(r)} \trianglelefteq \Gamma_r$  (*id* on a nbhd of  $r$ ).

**Step 1:** Each  $\Gamma_r$  acts with a global fixed point.

- Any circle action of  $\Gamma_{(r)}$  has a fixed point ( $H_b^2(\Gamma_{(r)}) = 0$ ).  
 uniformly perfect (suspension trick)  $g = a_g(ha_g^{-1}h^{-1})$   
 $H^k(\Gamma_{(r)}) = 0$  (Mather's suspension argument)



$$a_g = g \cdot (hgh^{-1}) \cdot (h^2gh^{-2}) \cdots$$

# Step 1 details/structural results

$r$  short ray,  $\Gamma_r := \text{Stab}(r)$ ,  $\Gamma_{(r)} \trianglelefteq \Gamma_r$  (*id* on a nbhd of  $r$ ).

**Step 1:** Each  $\Gamma_r$  acts with a global fixed point.

- Any circle action of  $\Gamma_{(r)}$  has a fixed point ( $H_b^2(\Gamma_{(r)}) = 0$ ).
- $\Gamma_r = \langle \Gamma_{(r)}, \Gamma_{(s)} \cap \Gamma_r \rangle$  if  $\text{end}(r) \neq \text{end}(s)$ .  
 $\Gamma_{(s)} \cap \Gamma_r$  preserves  $\text{Fix}(\Gamma_{(r)})$  and has fixed points.
- $\Gamma = \langle \Gamma_r, \Gamma_s \rangle$  if  $\text{end}(r) \neq \text{end}(s)$ .

**Cor:**  $\Gamma$  is generated by elements supported in disks.

use this + suspension to prove Theorem 2.



# Step 2 details

**Step 2:**  $P$  preserves the circular order and is injective.

- $P(r) \neq P(s)$  if  $\text{end}(r) \neq \text{end}(s)$  since  $\Gamma = \langle \Gamma_r, \Gamma_s \rangle$ .
- $\text{Or}(P(r_1), P(r_2), P(r_3)) = 1$  if  $\text{Or}(r_1, r_2, r_3) = 1$  and  $r_1, r_2, r_3$  are disjoint with distinct endpoints.

**goal:** remove “disjoint”.

# The filtration

Induction using a **filtration** associated to an “equator”  $\gamma$ .

**equator**: embedded circle containing  $K$ .

$R_0(\gamma) \subset R_1(\gamma) \subset \cdots \subset R$ , each is a Cantor set

each step adds Cantor to each complementary interval

# The induction

